



EPA REGISTER
HSR 100494
HSNO Classifications
6.1B (oral), 6.9A, 9.3A,
9.1 D

Registered pursuant to
the ACVM Act 1997
No. V9571
See
<https://eatsafe.nzfsa.govt.nz> for registration
conditions.

BATCH NO.
MANUFACTURED
DATE:
EXPIRY DATE:

NET CONTENTS: 22g
as 10 syringes each
containing 2.2g.

BAR CODE

DANGER

Keep out of reach of children

Read label before use

PredaSTOP™

for Feral Cats

***A single feed bait for the control of feral cats
(Felis catus)***

Contains 410 g/kg para-aminopropiophenone in the form
of a paste.

WARNING: May be fatal if swallowed in large quantities.
Reduces affinity of red blood cells for oxygen Very toxic
to some terrestrial vertebrates.

PRECAUTIONS: To be under control of Approved
Handler and holder of Controlled Substance Licence
(CSL). Wash hands thoroughly after handling. Do not eat,
drink or smoke when using this product. Prevent access
to bait by non-target domestic animals, wildlife and
livestock. Particular care should be taken if non-target
animals that are likely to be attracted to mince baits may
have access to the area being targeted (e.g. domestic
cats, dogs, kiwi and weka).

DIRECTIONS FOR USE: Use only in bait stations. Pre-
feeding before laying bait must be undertaken. Pre-
feeding should be continued until bait acceptance has
been established. Remove all pre-feed when laying toxic
bait. Follow Bait Preparation Instructions (enclosed and
available from www.connovation.co.nz). Meat baits
should be used within 48 hours of preparation and kept
as cool as possible prior to placing in stations.
Remove all pre-feed when laying toxic bait.
Bait take from pre-feeding should be used to determine
how many toxic mince baits to place in each station.
Where practical, ensuring that baits remain fresh may
maximise feral cat control.

On completion of the control operation, recover and bury
all remaining baits.

SYMPTOMS OF POISONING: Symptoms skin colour
changes (pale or blue-ish) lethargy and lack of co-
ordination.



	<p>FIRST AID: In the event of poisoning or suspected poisoning act immediately. Have product label or Safety Data Sheet available. Get medical attention. For advice contact National Poisons Centre 0800 POISON (0800 764 766). For animals, consult a veterinarian immediately. Product contains an anti-emetic. Administration of methylene blue antidote may be an option for non-target animals.</p>
--	--

STORAGE: Store locked up in original packaging tightly closed until used. Any product held after the expiry date shown on the container should be disposed of according to the disposal directions.

SPILLAGE: In the event of a spill, contain and isolate the spill and take all practical steps to manage any harmful effects. Scoop up spilled bait into secure labelled containers.

DISPOSAL: Dispose of unwanted or expired product either by burying under at least 60 cm of damp biologically active soil or transfer to an approved disposal facility. DO NOT use the empty container for any other purpose. Dispose of container by transfer to an approved disposal facility.

LEGAL OBLIGATIONS:

This product must only used in accordance with the label. Users must also meet the mandatory requirements of ACVM conditions of registration and HSNO controls (www.epa.govt.nz), including notification and signage requirements.

MANUFACTURED AND DISTRIBUTED BY: CONNOVATION LIMITED, 36B SIR WILLIAM DRIVE, EAST TAMAKI, AUCKLAND.
PO Box 58 613, BOTANY, AUCKLAND 2163, TEL (09) 273 4333
www.connovation.co.nz

